

Particle Collision near 1+1- Dimensional Horava-Lifshitz Black Hole and Naked Singularity

M. Halilsoy* and A. Ovgun†

Physics Department, Eastern Mediterranean University, Gazimagusa, TRNC, Mersin 10, Turkey.

(Dated: December 30, 2016)

The unbounded center of mass (CM) energy of oppositely moving colliding particles near horizon emerge also in 1+1- dimensional Horava-Lifshitz gravity. This theory has imprints of renormalizable quantum gravity characteristics in accordance with the method of simple power counting. Surprisingly the result obtained is not valid for a 1- dimensional Compton- like process between an outgoing photon and an infalling massless/ massive particle. It is possible to achieve unbounded CM energy due to collision between infalling photons and particles. The source of outgoing particles may be attributed to an explosive process just outside the horizon for a black hole and the naturally repulsive character for the case of a naked singularity. It is found that absence of angular momenta in 1+1- dimensions does not yield unbounded energy for collisions in the vicinity of naked singularities.

PACS numbers:

Keywords: Particle collision, BSW effect, Particle accelerators, Center-of-mass energy, Horava-Lifshitz gravity, Naked singularity

I. INTRODUCTION

It is known that in spacetime dimensions less than four gravity has no life of its own unless supplemented by external sources. With that addition we can have lower dimensional gravity and we can talk of black holes, wormholes, geodesics, lensing effect etc. in analogy with the higher dimensions. One effect that attracted much interest in recent times is the process of particle collisions near the horizon of black holes due to Banados, Silk and West [1] which came to be known as the BSW effect. This problem arose as a result of imitating the rather expensive venture of high energy particle collisions in laboratory. From curiosity the natural question arises: is there a natural laboratory (a particle accelerator) in our cosmos that we may extract information/ energy in a cheaper way? This automatically drew attentions to the strong gravity regions such as near horizon of black holes. Rotating black holes host greater energy reservoir due to their angular momenta and attentions naturally focused therein first [2, 3]. In case the metric is static and diagonal, there are reasons to consider the collision process in the vicinity of a naked singularity as well.

We note from physical grounds that outgoing particles from the event horizon of a black hole can not occur. Hawking radiation particles/ photons emerge too weak to compare with infalling particles. Thus collision of two particles can only be argued if both are infalling toward the horizon of a black hole. Such a process, however, yield no BSW effect in the non- rotating metrics, which is our main interest in this study. In order to have an unbounded CM energy in a collision process both particles must be taken in the same coordinate frame and in opposite directions. This is possible in the vicinity of a naked singularity whose repulsive effect compells particles/ photons to make collisions with an infalling particle/ photon. From the outset we state that such a collision taking place near the naked singularity in the absence of angular momenta does not yield an unbounded CM energy. To extend our study to cover also collisions near black holes we assume that some unspecified process, such as disintegration decay process of some particles, yield outgoing particles photons while the partners fall into the hole. For a thorough analysis of all these problems covering the ergosphere region of a Kerr black hole, Penrose process, particle collisions etc. one must consult [48].

In general one considers the radial geodesics and upon energy-momentum conservation in the center-of-mass (CM) frame the near horizon limit is checked whether the energy is bounded/unbounded. Our aim in this study is to consider black hole solutions in 1+1- dimensional Hořava-Lifshitz (HL) gravity [4] and check the BSW effect in such reduced dimensional theory. Let us remark that at the Planck scale in higher dimensions the spherical part $r^2 d\Omega_{D-2}^2$ of the line element is less effective compared to the time and radial components. For this reason 1+1- dimensions becomes significant at the Planck scale. For a number of reasons HL gravity is promising as a candidate for a re-normalizable quantum gravity physics which has been yearned for a long time [5]. The key idea in HL- gravity is the inhomogeneous scaling properties of time and space coordinates which violates the Lorentz invariance. Arnowitt-Deser-Misner (ADM)

*Electronic address: mustafa.halilsoy@emu.edu.tr

†Electronic address: ali.ovgun@emu.edu.tr

splitting of space and time[6] constitutes its geometrical background. BSW effect in lower/higher- dimensions has been worked out by many authors [7–43]. Following the similar idea we consider black hole solutions and naked singularities in 1+1- dimensions and search for the same effect in this lower dimension. It should be added that with 1+1- dimensional HL theory the simplest non-trivial solution is the solution describing an accelerated particle in the flat space of Rindler frame. This justifies also the meaning of the vector field (a_i) as the acceleration in the HL-gravity. The role of Rindler acceleration in 3+1- dimensions as a possible source of flat rotation curves and geodesics motion has been discussed recently [44]. It is our belief that the results in lower dimensions are informative for higher dimensions and as a toy model can play the role as precursors in this regard. Even a Compton- like process can be considered at the toy level between a massless photon outgoing from a naked singularity and a particle falling into the naked singularity. The diverging CM energy results in the case of photon-particle collision in 1+1- dimensions under specific conditions.

Organization of the paper as follows. In Section II, we review in brief the 1+1-D HL theory with a large class of black hole and naked singularity solutions. CM energy of colliding particles near horizon and naked singularity is considered in Section III. Section IV proceeds with applications to particular examples. The case of particle-photon collision is studied separately in Section V. The paper ends with our conclusion in Section VI.

II. 1+1-D HL BLACK HOLE/ NAKED SINGULARITY

HL formalism in 3+1-D makes use of the ADM splitting of time and space components as follows

$$ds^2 = -N^2 dt^2 + g_{ij} (dx^i + N^i dt) (dx^j + N^j dt) \quad (1)$$

where $N(t)$ and N^i are the lapse and shift functions, respectively. The action of this theory is

$$S = \frac{M_{Pl}^2}{2} \int d^3x dt \sqrt{g} (K_{ij} K^{ij} + \lambda K^2 + V(\phi)) \quad (2)$$

where K_{ij} is the extrinsic curvature tensor with trace K and Planck mass M_{Pl} . $V(\phi)$ stands for the potential function of a scalar field ϕ , and λ is a constant ($\lambda > 1$). Reduction from 3+1- D to 1+1- D results in the action [4]

$$S = \int dt dx \left(-\frac{1}{2} \eta N^2 a_1^2 + \alpha N^2 \phi'^2 - V(\phi) \right) \quad (3)$$

where η =constant and α =constant will be chosen to be unity and $a_1=(\ln N)'$. Let us comment that a ‘prime’ denotes $\frac{d}{dx}$. We note that also the first term in S is inherited from the geometric part of the action while the other two terms are from the scalar field source. For simplicity we have set also $M_{Pl} = 1$.

It has been shown in [4] that by variational principle a general class of solutions is obtained as follows

$$N(x)^2 = 2C_2 + \frac{A}{\eta} x^2 - 2C_1 x + \frac{B}{\eta x} + \frac{C}{3\eta x^2} \quad (4)$$

in which C_2 , A , C_1 , B and C are integration constants. Ref.[4] must be consulted for the physical content of these constants.

The line element is

$$ds^2 = -N(x)^2 dt^2 + \frac{dx^2}{N(x)^2} \quad (5)$$

with the scalar field

$$\phi(x) = \ln \sqrt{2C_2 + \frac{A}{\eta} x^2 - 2C_1 x + \frac{B}{\eta x} + \frac{C}{3\eta x^2}}. \quad (6)$$

Note that the associated potential is

$$V(\phi(x)) = A + \frac{B}{x^3} + \frac{C}{x^4}. \quad (7)$$

and the Ricci scalar is calculated as

$$R = -\frac{2}{\eta} \left(A + \frac{B}{x^3} + \frac{C}{x^4} \right). \quad (8)$$

There is naked singularity when $A = C_1 = 0$ and $C_2 = B = C = \eta = 1$, so that there is no horizon for

$$N(x)^2 = 2 + \frac{1}{x} + \frac{1}{3x^2}. \quad (9)$$

Another black hole solution reported by Bazeia et. al. [4] is found by taking $C_1 \neq 0$, $C_2 \neq 0$, $B \neq 0$ and $A = C = 0$

$$N(x)^2 = 2C_2 - 2C_1x + \frac{B}{\eta x}. \quad (10)$$

This solution develops the following horizons

$$x_h^\pm = \frac{C_2}{2C_1} \pm \sqrt{\Delta}, \quad \Delta = \frac{C_2^2}{4C_1^2} + \frac{B}{2\eta C_1}. \quad (11)$$

As $\Delta = 0$ they degenerate, i.e., $x_h^+ = x_h^-$.

The Hawking temperature is given in terms of the outer (x_h^+) horizon as follows

$$T_H = \frac{(N(x)^2)'}{4\pi} \bigg|_{x=x_h^+}. \quad (12)$$

For the special case $C_2 = 0$, $C_1 = -M$ and $B = -2M$ the horizons are independent of the mass M :

$$x_h^\pm = \pm \frac{1}{\sqrt{\eta}} \quad (\eta > 0) \quad (13)$$

The temperature is then given simply by

$$T_H = \frac{M}{\pi}. \quad (14)$$

This is a typical relation between the Hawking temperature and the mass of black holes in 1 + 1 dimensions [45].

In the case of $C_2 = 1/2$, $B = -2M$, $\eta = 1$ and $A = C = C_1 = 0$, it gives a Schwarzschild-like solution;

$$N(x)^2 = 1 - \frac{2M}{x} \quad (15)$$

On the other hand, the choice of the parameters, for $C_2 = 1/2$, $B = -2M$, $C = 3Q^2$, $\eta = 1$ and $A = C_1 = 0$ gives a Reissner–Nordström-like solution.

$$N(x)^2 = 1 - \frac{2M}{x} + \frac{Q^2}{x^2} \quad (16)$$

As in the general relativity we can make particular choice of the parameters so that we end up with a naked singularity instead of a black hole. The choice $Q^2 > M^2$ in eq.(16), for instance, transforms the HL- black hole into a naked singularity at $x = 0$. Similarly $M < 0$, turns eq.(15) into a naked singular metric at $x = 0$.

III. CM ENERGY OF PARTICLE COLLISION NEAR THE HORIZON OF THE 1+1 -D HL BLACK HOLE

Here we will derive the equations of motion of an uncharged massive test particle by using the method of geodesic Lagrangian. Such equations can be derived from the Lagrangian equation,

$$\mathcal{L} = \frac{1}{2} [-N(x)^2 \left(\frac{dt}{d\tau} \right)^2 + \frac{1}{N(x)^2} \left(\frac{dx}{d\tau} \right)^2] \quad (17)$$

in which, τ is the proper time for time-like geodesics (or massive particles). The canonical momenta are

$$p_t = \frac{d\mathcal{L}}{dt} = -N(x)^2 \dot{t} \quad (18)$$

and

$$p_x = \frac{d\mathcal{L}}{dx} = \frac{\dot{x}}{N(x)^2} \quad (19)$$

The 1+1- D HL black hole have only one Killing vector ∂_t . The associated conserved quantity will be labeled by E . From eq.(18), E is related to $N(x)^2$ as,

$$-N(x)^2 \dot{t} = -E \quad (20)$$

Hence,

$$\dot{t} = \frac{E}{N(x)^2} \quad (21)$$

The two-velocity of the particles are given by $u^\mu = \frac{dx^\mu}{d\tau}$. We have already obtained u^t in the above derivation. To find $u^x = \dot{x}$, the normalization condition for time-like particles, $u^\mu u_\mu = -1$ [1, 46] can be used as,

$$g_{tt}(u^t)^2 + g_{xx}(u^x)^2 = -1 \quad (22)$$

By substituting u^t to eq.(22), one obtain u^x as,

$$(u^x)^2 = E^2 - N(x)^2 \quad (23)$$

for which an effective potential V_{eff} can be defined by

$$(u^x)^2 + V_{eff} = E^2. \quad (24)$$

Now, the two-velocities can be written as,

$$u^t = \dot{t} = \frac{E}{N(x)^2} \quad (25)$$

and

$$u^x = \dot{x} = \sqrt{E^2 - N(x)^2}. \quad (26)$$

We proceed now to present the CM energy of two particles with two-velocities u_1^μ and u_2^μ . We will assume that both have rest mass $m_0 = 1$. The CM energy is given by,

$$E_{cm} = \sqrt{2} \sqrt{(1 - g_{\mu\nu} u_1^\mu u_2^\nu)} \quad (27)$$

so

$$\frac{E_{cm}^2}{2} = 1 + \frac{E_1 E_2}{N(x)^2} - \frac{\kappa \sqrt{E_1^2 - N(x)^2} \sqrt{E_2^2 - N(x)^2}}{N(x)^2} \quad (28)$$

where $\kappa = \pm 1$ corresponds to particles moving in the same/ opposite direction with respect to each other. We wish to stress that our concern is for the case $\kappa = \pm 1$ since no physical particle is ejected from the black hole. Note that E_1 and E_2 are the energy constants corresponding to each particle. In the case the second term under the square root is too small than the first one,

$$\sqrt{E^2 - N(x)^2} \approx \left(E - \frac{N(x)^2}{2E^2} + \dots \right) \quad (29)$$

so that the higher order terms can be neglected and CM energy of two particles can be written as [22]

$$\frac{E_{cm}^2}{2} \approx 1 + (1 - \kappa) \frac{E_1 E_2}{N(x)^2} + \frac{\kappa}{2} \left(\frac{E_2}{E_1} + \frac{E_1}{E_2} \right) \quad (30)$$

Obviously the case with $\kappa = +1$, in which the CM energy becomes

$$\frac{E_{cm}^2}{2} \approx 1 + \frac{(E_2^2 + E_1^2)}{2E_1E_2}, \quad (31)$$

where the CM energy is independent from metric function, and it gives always a finite energy. On the other hand $\kappa = -1$ gives

$$\frac{E_{cm}^2}{2} \approx 1 + \frac{2E_1E_2}{N(x)^2} - \frac{(E_2^2 + E_1^2)}{2E_1E_2} \quad (32)$$

in which it gives unbounded CM energy near the horizon of the HL black holes provided an outgoing particle mechanism from the horizon is established. Otherwise the yield of two ingoing particles collision remains finite.

IV. SOME EXAMPLES

A. Schwarzschild-like Solution

In the case of $C_2 = 1/2$, $B = -2M$, $\eta = 1$ and $A = C = C_1 = 0$, it gives Schwarzschild-like solution where

$$V(\phi(x)) = -\frac{2M}{x^3} \quad (33)$$

and

$$N(x)^2 = 1 - \frac{2M}{x} \quad (34)$$

For the CM energy on the horizon, we have to compute the limiting value of eq.(28) as $x \rightarrow x_h = 2M$, where is the horizon of the black hole. Setting $\kappa = -1$ as is, the CM energy near the event horizon for 1+1 D Schwarzschild BH is

$$E_{cm}^2(x \rightarrow x_h) = \infty \quad (35)$$

This result for 4-D Schwarzschild Black hole is already calculated by Baushev [23]. Hence, the condition of $\kappa = -1$, when the location of particle 1 approaches the horizon, on the other hand the particle 2 runs outward from the horizon due to some unspecified physical process yet yields $E_{cm}^2 \rightarrow \infty$ so there is BSW effect for 1+1 Schwarzschild-like Solution when the condition $\kappa = -1$ is satisfied.

B. Reissner-Nordstrom-like solution

On the other hand, the choice of the parameters, for $C_2 = 1/2$, $B = -2M$, $C = 3Q^2$, $\eta = 1$ and $A = C_1 = 0$ gives the Reissner–Nordström-like solution.

$$N(x)^2 = 1 - \frac{2M}{x} + \frac{Q^2}{x^2} \quad (36)$$

and

$$V(\phi(x)) = -\frac{2M}{x^3} + \frac{3Q^2}{x^4} \quad (37)$$

so the CM energy is calculated by using the limiting value of eq.(32)

$$E_{cm}^2(x \rightarrow x_{h=M+\sqrt{M^2-Q^2}}) = \infty \quad (38)$$

so there is a BSW effect.

C. The Extremal case of the Reissner-Nordstrom like black hole

For the extremal case we have with $M = Q$, from eq.(36)

$$N(x)^2 = \left(1 - \frac{M}{x}\right)^2 \quad (39)$$

so that it also gives the same answer from eq.(32) as

$$E_{cm}^2(x \rightarrow x_h) = \infty. \quad (40)$$

D. Specific New Black Hole Case

The new 3-parametric black hole solution given by Bazeia, Brito and Costa [4] is chosen as

$$N(x)^2 = 2C_2 - 2C_1x + \frac{B}{\eta x} \quad (41)$$

with the potential

$$V(\phi(x)) = \frac{B}{x^3}. \quad (42)$$

For the special case $C_2 = 0$, $C_1 = -M$ and $B = -2M$ we have

$$N(x)^2 = 2Mx - \frac{2M}{\eta x} \quad (43)$$

with suitable potential which is

$$V(\phi(x)) = -\frac{2M}{x^3}. \quad (44)$$

The CM energy of two colliding particles is calculated by taking the limiting values of eq.(32)

$$E_{cm}^2(x \rightarrow x_h) = \infty \quad (45)$$

Hence the BSW effect arises here as well.

E. Near Horizon Coordinates

We have explored the region near the horizon by replacing r by a coordinate ρ . The proper distance from the horizon ρ [47]:

$$\rho = \int \sqrt{g_{xx}(x')} dx' = \int_{x_h}^x \frac{1}{N(x')} dx' \quad (46)$$

The first example is the Schwarzschild-like solution which is

$$N(x)^2 = 1 - \frac{2M}{x} \quad (47)$$

so that proper distance is calculated as

$$\begin{aligned} \rho &= \int_{x_h}^x \left(1 - \frac{2M}{x'}\right)^{-\frac{1}{2}} dx' \\ &= \sqrt{x(x-2M)} + 2MG \sinh^{-1}(\sqrt{x/2M-1}). \end{aligned} \quad (48)$$

The new metric is

$$ds^2 = -\left(1 - \frac{2M}{x(\tilde{\rho})}\right)dt^2 + d\tilde{\rho}^2 \quad (49)$$

where $\tilde{\rho} \simeq 2\sqrt{2M(x-2M)}$ so that it gives approximately

$$ds^2 \simeq -\frac{\rho^2}{(4M)^2}dt^2 + d\rho^2 \quad (50)$$

which is once more the Rindler type line element. Let us note that this Rindler- type line element is valid within the near horizon limit approximation. For practical purposes there are advantages in adapting such an approximation which is also in conform with the equivalence principle [47]. The CM energy of two colliding particles is given by

$$\frac{E_{cm}^2}{2m_0^2} = 1 + \frac{(4M)^2 \left(E_1 E_2 - \kappa \sqrt{E_1^2 - \frac{\rho^4}{(4M)^4}} \sqrt{E_2^2 - \frac{\rho^4}{(4M)^4}} \right)}{\rho^2} \quad (51)$$

so that there is BSW effect for $\kappa = -1$ when $\rho \rightarrow 0$.

V. PARTICLE COLLISION NEAR THE NAKED SINGULARITY

There is a naked singularity for our 1+1- HL model at the location of $x = 0$, with $Q^2 > M^2$ in eq.(16). In addition $M < 0$, turns eq.(15) into a naked singular metric at $x = 0$. There is also naked singularity when we choose metric function as follows

$$N(x)^2 = 2 + \frac{1}{x} + \frac{1}{3x^2} = \frac{6x^2 + 3x + 1}{3x^2}. \quad (52)$$

As it is given in eq.(28), CM energy of the collision of two particles generally is (for $N(x) \rightarrow \infty$)

$$\frac{E_{cm}^2}{2} \approx 1 - \kappa + \frac{1}{2N(x)^2} [2E_1 E_2 + \kappa(E_1^2 + E_2^2)]. \quad (53)$$

For the case $\kappa = \pm 1$, when x goes to zero, the CM energy remains finite for radially moving particles

$$\frac{E_{c.m.}^2}{2} \Big|_{x=0} \rightarrow 1 - \kappa. \quad (54)$$

This suggests that although one of the particle is boosted by the naked singularity. There is not any unlimited collisional energy near such a singularity. Note that Compton- like processes were considered first in [48], where rotational effect of Kerr black hole played a significant role. Our case here is entirely free of rotational effects.

VI. PHOTON VERSUS AN INFALLING PARTICLE

A massless photon can naturally scatter an infalling particle or vice versa. This phenomenon is analogous to a Compton scattering taking place in 1+1-dimensions. Null-geodesics for a photon can be described simply by

$$\begin{aligned} \frac{dt}{d\lambda} &= \frac{E_1}{N^2} \\ \frac{dx}{d\lambda} &= \pm \sqrt{E_1^2 - N^2} \end{aligned} \quad (55)$$

where λ is an affine parameter and E_1 stands for the photon energy. Defining $E_1 = \hbar\omega_0$, where ω_0 is the frequency (with the choice $\hbar = 1$) we can parametrize energy of the photon by ω_0 alone. The CM energy of a photon and the infalling particle can be taken now as

$$E_{cm}^2 = -(p^\mu + k^\mu)^2 \quad (56)$$

in which $p^\mu = mu^\mu$ and k^μ refer to the particle and photon, 2- momenta, respectively. This amounts to

$$E_{cm}^2 = m^2 - 2mg_{\mu\nu}u^\mu k^\nu, \quad (57)$$

where we have for the particle

$$p^\mu = m \left(\frac{E_2}{N^2}, \sqrt{E_2^2 - N^2} \right) \quad (58)$$

and for the photon

$$k^\mu = \left(\frac{E_1}{N^2}, -E_1 \right). \quad (59)$$

One obtains

$$E_{cm}^2 = m^2 + \frac{2mE_1}{N^2} \left(E_2 + \kappa \sqrt{E_2^2 - N^2} \right). \quad (60)$$

In the near horizon limit this reduces to

$$E_{cm}^2 = m^2 + \frac{2mE_1}{N^2} \left(E_2 + \kappa E_2 - \frac{N^2}{2E_2} \right). \quad (61)$$

Note that for $\kappa = -1$ we have E_{cm}^2 given by

$$E_{cm}^2 = m^2 \left(1 - \frac{E_1}{mE_2} \right) \quad (62)$$

which is finite between the collision of a photon and an infalling particle, and therefore is not of interest. As a matter of fact the occurrence of outgoing photon from the event horizon can not be justified unless an explosive/decay process is assumed to take place. As a result for $\kappa = +1$ from eq.(61) we obtain an unbounded E_{cm}^2 between the collision of infalling photon and particle. Let us add that ‘inverse’ Compton process in the ergosphere of Kerr black hole was considered in [48] where the photon’s energy showed increment due to rotational and curvature effects. The energy, however, attained an upper bound which was finite. Our result obtained here being entirely radial on the other hand can hardly be compared with those of [48].

VII. CONCLUSION

Our aim was to investigate whether the BSW type effect which arises in higher dimensional black holes applies also in the 1+1- D naked singularity/ black hole. The theory we adapted is not general relativity but instead the recently popular HL gravity. We employed the class of 5- parametric black hole/ naked singularity solutions found recently [4]. The class has particular limits of flat Rindler, Schwarzschild and Reissner-Nordstrom- like solutions. For each case we have calculated the center-of-mass (CM) energy of the particles and shown that the energy can grow unbounded for some cases. In other words the strong gravity near the event horizon effects the collision process with unlimited source to turn it into a natural accelerator. The model we use applies also to the case of a photon/particle collision with different characteristics. It is observed that the CM energy of the infalling particles from rest at infinity will remain finite in the CM frame at the event horizon of a black hole. Contrariwise, unlimited CM energy will be attained between the collision of the outgoing particles from the event horizon region and infalling particles. It is also possible to achieve the infinite energy between an infalling photon and an infalling massive particle. However, we found finite CM energy between an outgoing photon and infalling particle. Finally, we must admit that absence of rotational effects in 1+1- D restricts the problem to the level of a toy model in which particles move on pure radial geodesics yielding finite CM energy in the vicinity of a naked singularity.

VIII. ACKNOWLEDGEMENT

We wish to thank the anonymous referees for valuable suggestions. This work was presented as a poster at Karl Schwarzschild Meeting 20-24 July 2015 Frankfurt Institute for Advanced Studies.

- [2] Ted Jacobson, Thomas P. Sotiriou, Phys. Rev. Lett. **104** (2010), 021101.
- [3] Kayll Lake, Phys. Rev. Lett. **104** (2010), 211102.
- [4] D. Bazeia, F. A. Brito, F. G. Costa, Phys. Rev. D. **91** (2015), 044026.
- [5] P. Horava, Phys. Rev. D. **79** (2009), 084008.
- [6] R. Arnowitt, S. Deser, C. W. Misner, Gen. Relativ. Gravit. **40** (2008), 1997.
- [7] E. Berti, V. Cardoso, L. Gualtieri, F. Pretorius and U. Sperhake, Phys. Rev. Lett. **103** (2009), 239001.
- [8] M. Ban ados, B. Hassanain, J. Silk, S. M. West, Phys. Rev. D **83** (2011), 023004.
- [9] T. Jacobson, T. P. Sotiriou, Phys. Rev. Lett. **104** (2010), 021101.
- [10] O. B. Zaslavskii, JETP Lett. **92** (2010), 571.
- [11] S. W. Wei, Y. X. Liu, H. T. Li, F. W. Chen, J. High Energy Phys. **12** (2010), 066.
- [12] O. B. Zaslavskii, Phys. Rev. D. **86** (2012), 124039.
- [13] H. Saadat, Can. J. Phys. (2014) **92** (12):1562.
- [14] N. Tsukamoto, C. Bambi, Phys. Rev. D. **91** (2015), 084013.
- [15] S. G. Ghosh, P. Sheoran, M. Amir, Phys. Rev. D. **90** (2014), 103006.
- [16] A. Galajinsky, Phys. Rev. D. **88** (2013), 027505.
- [17] V. P. Frolov, Phys. Rev. D. **85** (2012), 024020.
- [18] A. Al Zahrani, V. P. Frolov, A. Shoom, Phys. Rev. D. **87** (2013), 084043.
- [19] J. Sadeghi, B. Pourhassan, Eur. Phys. J. C **72** (2012), 1984.
- [20] J. Sadeghi, B. Pourhassan and H. Farahani, Commun. Theor. Phys. **62** (2014), 358-362.
- [21] C. Liu, S. Chen, C. Ding, J. Jian, Phys. Lett. B **701** (2011), 285.
- [22] M. Patil, P. S. Joshi, Phys. Rev. D. **86** (2012), 044040.
- [23] A. N. Baushev, Int. J. Mod. Phys. D. **18** (2009), 1195-1203.
- [24] M. Patil, P. S. Joshi, Gen. Relativ. Gravit. **46** (2014), 1801.
- [25] J. D. Schnittman, Phys. Rev. Lett. **113** (2014), 261102.
- [26] M. Patil, P. S. Joshi, Phys. Rev. D **82** (2010), 104049.
- [27] M. Patil, P. S. Joshi and D. Malafarina, Phys. Rev. D **83** (2011), 064007.
- [28] A. A. Grib, Yu. V. Pavlov, Astropart. Phys. **34** (2011), 581.
- [29] M. Sharif, N. Haider, Astrophys. Space Sci. **346** (2013), 111-117.
- [30] I. Hussain, M. Jamil, B. Majeed, Int. J. Theor. Phys. **54** (2015) 5, 1567-1577.
- [31] S. Hussain, I. Hussain, M. Jamil, Eur. Phys. J. C **74** (2014) 12, 3210.
- [32] M. Amir, S. G. Ghosh, JHEP 1507 (2015), 015.
- [33] B. Pourhassan, U. Debnath, arXiv:1506.03443.
- [34] A. A. Grib, Yu. V. Pavlov, Grav. Cosmol. **21** (2015), 13-18.
- [35] A. A. Grib, Yu. V. Pavlov, Theor. Math. Phys. **185** (2015) 1, 1425-1432.
- [36] C. Ding, C. Liu, Q. Guo, Int. J. Mod. Phys. D **22** (2013), 1350013.
- [37] Jie Yang, Yun-Liang Li, Yang Li, Shao-Wen Wei, Yu-Xiao Liu, Adv. High Energy Phys. **2014** (2014), 204016.
- [38] H. Nemoto, U. Miyamoto, T. Harada, T. Kokubu, Phys. Rev. D **87** (2013) 12, 127502.
- [39] C. Zhong, S. Gao, JETP Lett. **94** (2011), 589-592.
- [40] C. Liu, S. Chen, J. Jing, Chin. Phys. Lett. **30** (2013), 100401.
- [41] Yi Zhu, Shao-Feng Wu, Yu-Xiao Liu, Ying Jiang, Phys. Rev. D **84** (2011), 043006.
- [42] U. Miyamoto, H. Nemoto, M. Shimano, Phys. Rev. D **83** (2011), 084054.
- [43] Yang Li, Jie Yang, Yun-Liang Li, Shao-Wen Wei, Yu-Xiao Liu, Class. Quant. Grav. **28** (2011), 225006.
- [44] M. Halilsoy, O. Gurtug, S. Habib Mazharimousavi, Gen. Rel. Grav. **45** (2013), 2363-2381.
- [45] S. W. Hawking, Phys. Rev. D. **13** (1976), 191.
- [46] C. W. Misner, K. S. Thorne and J. A. Wheeler, San Francisco 1973, 1279p.
- [47] L. Susskind and J. Lindesay, Hackensack, USA: World Scientific (2005) 183 p.
- [48] T. Piran and J. Shaham, Phys. Rev. D. **16**, 6 (1977).